arXiv:1602.04866v1 [math-ph] 16 Feb 2016

A FERMI GOLDEN RULE FOR QUANTUM GRAPHS

MINJAE LEE AND MACIEJ ZWORSKI

Abstract. We present a Fermi golden rule giving rates of decay of states obtained by perturbing embedded eigenvalues of a quantum graph. To illustrate the procedure in a notationally simpler setting we also present a Fermi Golden Rule for boundary value problems on surfaces with constant curvature cusps. We also provide a resonance existence result which is uniform on compact sets of energies and metric graphs. The results are illustrated by numerical experiments.

1. Introduction and statement of results

Quantum graphs are a useful model for spectral properties of complex systems. The complexity is captured by the graph but analytic aspects remain one dimensional and hence relatively simple. We refer to the monograph by Berkolaiko–Kuchment [1] for references to the rich literature on the subject.

In this note we are interested in graphs with infinite leads and consequently with continuous spectra. We study dissolution of embedded eigenvalues into the continuum and existence of resonances close to the continuum. Our motivation comes from a recent Physical Review Letter [10] by Gnutzmann–Schanz–Smilansky and from a mathematical study by Exner–Lipovský [9].

We consider an oriented graph with vertices $\{v_j\}_{j=1}^J$, infinite leads $\{e_k\}_{k=1}^K$, $K>0$, and M finite edges $\{e_m\}_{m=K+1}^{M+K}$. We assume that each finite edge, e_m, has two distinct vertices as its boundary (a non-restrictive no-loop condition) and we write $v\in e_m$ for these two vertices v. An infinite lead has one vertex. The set of (at most two) common vertices of e_m and e_ℓ is denoted by $e_m\cap e_\ell$ and we we denote by $e_m\ni v$ the set of all edges having v as a vertex.

The finite edges are assigned length ℓ_m, $K+1\leq m\leq M+K$ and we put $\ell_k=\infty$, $1\leq k\leq K$, for the infinite edges. To obtain a *quantum graph* we define a Hilbert space, is given by

$$L^2:=\bigoplus_{m=1}^{K+M} L^2([0,\ell_m]),\quad L^2\ni u=(u_1,\cdots u_{M+K}),\quad u_m\in L^2([0,\ell_m]).$$



FIGURE 1. A graph given by a cycle $\{e_k\}_{k=K+1}^{2K}$ connected to K infinite leads $\{e_k\}_{k=1}^{K}$ at K vertices: v_k, $e_{K+k}\cap e_{K+k-1} = v_k$, $e_{2K}\cap e_{K+1} = v_1$, $e_k\cap e_{K+k} = v_k$. The lengths of finite edges are given by $\ell_k(t) = e^{-2a_k(t)}\ell_k$, $K+1\leq k\leq 2K$. If $\ell_k(0)$'s are rationally related then $P(0)$ has eigenvalues, $\lambda(0)$, embedded in the continuous spectrum. If $\lambda(0)$ is simple then $\lambda(0)$ belongs to a smooth family of resonances, $\lambda(t)$, $\operatorname{Im}\lambda(t)\leq 0$. Theorem 1 and Example 1 in §3 show that in this case $\operatorname{Im}\ddot\lambda = \lambda^2\sum_{k=1}^K|\langle\dot a u, e^k(\lambda)\rangle|^2$, where u is the normalized eigenfuction corresponding to u and $e^k(\lambda)$ is the generalized eigenfuction normalized in the kth lead – see (1.1).

We then consider the simplest quantum graph Hamiltonian which is unbounded operator P on L^2 defined by $(Pu)_m = -\partial_x^2 u_m$ with

$$\mathcal{D}(P) = \{u : u_m\in H^2([0,\ell_m]),\ u_m(v) = u_\ell(v),\ v\in e_m\cap e_\ell,\ \sum_{e_m\ni v}\partial_\nu u_m(v) = 0\}.$$

Here ∂_ν denotes the outward pointing normal at boundary of e_v:

$$u_m\in H^2([0,\ell_m]),\quad \partial_\nu u_m(0) = -u_m'(0),\quad \partial_\nu u_m(\ell_m) = u_m'(\ell_m).$$

Quantum graphs with infinite leads fit neatly into the general abstract framework of *black box* scattering [13] and hence we can quote general results [8, Chapter 4] in spectral and scattering theory.

When $K>0$ then the projection on the continuous spectrum of P is given in terms of generalized eigenfunctions $e^k(\lambda)$, $1\leq k\leq K$, which for $\lambda\notin \operatorname{Spec}_{\rm pp}(P)$ are characterized as follows:

$$\begin{gathered} e^k(\lambda)\in\mathcal{D}_{\rm loc}(P),\quad (P-\lambda^2)e^k(\lambda) = 0,\\ e^k_m(\lambda,x) = \delta_{mk}e^{-i\lambda x} + s_{mk}(\lambda)e^{i\lambda x},\quad 1\leq m\leq K.\end{gathered}\tag{1.1}$$

The family $\lambda\mapsto e^k(\lambda)\in\mathcal{D}_{\rm loc}(P)$ extends holomorphically to a neighbourhood of $\mathbb{R}$ and that defines $e^k(\lambda)$ for all λ. We will in fact be interested in $\lambda\in\operatorname{Spec}_{\rm pp}(P)$. The functions e^k parametrize the continuous spectrum of P – see [8, §4.4] and (3.13) below.

We now consider a family of quantum graphs obtained by varying the lengths ℓ_m, $K+1\leq m\leq M+K$:

$$\ell_m(t) = e^{-a_m(t)}\ell_m,\quad a_m(0) = 0.\tag{1.2}$$



FIGURE 2. A simple graph with embedded eigenvalues, $M = K = 2$. Solid lines and dashed lines indicate the trajectory of $\lambda(t)$ and of the second order approximation $\tilde{\lambda}(t) = \lambda + t\dot{\lambda} + \frac{i}{2}t^2 \operatorname{Im}\ddot{\lambda}$, respectively. (The colour coding indicates the parameter t shown in the colour bar.) We approximate the real part linearly using (3.11) and the imaginary quadratically using (1.3). The four cases are (a): $\ell_3(t) = 1 - t,\ \ell_4(t) = 1 - t$, (b): $\ell_3(t) = 1 - t,\ \ell_4(t) = 1$, (c): $\ell_3(t) = 1 - t,\ \ell_4(t) = 1 + t$, (d): $\ell_3(t) = 1 - t,\ \ell_4(t) = 1 + 2t$.

and the corresponding family of operators, $P(t)$. The works [9] and [10] considered the case in which $P(0)$ has embedded eigenvalues and investigated the resonances of the deformed family $P(t)$ converging to these eigenvalues as $t \to 0$. Here we present a Fermi golden rule type formula (see §2 for references to related mathematical work) which gives an infinitesimal condition for the disappearance of an embedded eigenvalue. It becomes a resonance of P and one can calculate the infinitesimal rate of decay. Resonances are defined as poles of the meromorphic continuation of $\lambda \mapsto (P - \lambda^2)^{-1}$ to $\mathbb{C}$ as an operator $L^2_{\text{comp}} \to L^2_{\text{loc}}$ (see [8, §4.2] and for a self-contained general argument Proposition 4.1). We denote the set of resonances of P by $\operatorname{Res}(P)$.

Theorem 1. *Suppose that $\lambda^2 > 0$ is a simple eigenvalue of $P = P(0)$ and u is the corresponding normalized eigenfunction. Then for $|t| \leq t_0$ there exists a smooth function $t \mapsto \lambda(t)$ such that $\lambda(t) \in \operatorname{Res}(P)$ and*

$$\operatorname{Im}\ddot{\lambda} = -\sum_{k=1}^{K} |F_k|^2,$$
$$F_k := \lambda\langle \dot{a}u, e^k(\lambda)\rangle + \lambda^{-1} \sum_{v} \sum_{e_m \ni v} \tfrac{1}{4}\dot{a}_m (3\partial_\nu u_m(v)\overline{e^k(\lambda, v)} - u(v)\partial_\nu \overline{e^k_m(\lambda, v)}) \tag{1.3}$$

The proof is given in §3 and that section is concluded with two examples: the first gives graphs and eigenvalues for which $F_k = \lambda\langle \dot{a}u, e^k(\lambda)\rangle$ – see Figures 1 and 2. The second example gives a graph and an eigenvalue for which the boundary terms in the formula for F_k are needed – see Fig. 4.

The formula (1.3) gives a condition for the existence a resonance with a nontrivial imaginary part (decay rate) near an embedded eigenvalue of the unperturbed operator: $D(\lambda_0, ct) \cap \mathrm{Res}(P(t)) \neq \emptyset$ for some c and for $|t| \leq t_0$, where the constants c and t_0 depend on λ_0 and $P(t)$. However, it is difficult to estimate the speed with which the resonance $\lambda(t)$ moves – that is already visible in comparing Fig. 2 with Fig. 4. (A striking example is given by $P(t) = -\partial_x^2 + tV(x)$ where $V \in C_c^\infty(\mathbb{R})$ and $t \to 0$; infinitely many resonances for $t \neq 0$ [20] disappear and $P(0)$ has only one resonance at 0.) Also, the result is not uniform if we vary λ_0 or the lengths of the edges.

The next theorem adapts the method of Tang–Zworski [18] and Stefanov [15] (see also [8, §7.3]) to obtain existence of resonances near any approximate eigenvalue and in particular near an embedded eigenvalue – see the example following the statement. In particular this applies to the resonances studied in [9] and [10]. The method applies however to very general Hamiltonians – for semiclassical operators on graphs the general black box resuls of [18] and [15] apply verbatim. The point here is that the constants are uniform even though the dependence on t is slightly weaker.

To formulate the result we define $D(\lambda_0, r) = \{\lambda \in \mathbb{C} : |\lambda - \lambda_0| < r\}$ and

$$\mathcal{H}_R := \bigoplus_{m=1}^{K} L^2([0, R]) \oplus \bigoplus_{m=K+1}^{K+M} L^2([0, \ell_m]). \tag{1.4}$$

Theorem 2. *Suppose that P is defined above and the lengths, ℓ_m, have the property that $\ell_m \in \mathcal{L}$, $K + 1 \leq m \leq M + K$ where $\mathcal{L}$ is a fixed compact subset of the the open half-line.*

Then for any $\mathcal{L} \Subset (0, \infty)$, $I \Subset (0, \infty)$, $R > 0$ and $\gamma < 1$ there exists $\varepsilon_0 > 0$ such that

$$\exists\, u \in \mathcal{H}_R \cap \mathcal{D}(P),\ \lambda_0 \in I \quad \text{such that} \quad \|u\|_{L^2} = 1, \quad \|(P - \lambda_0^2)u\| = \varepsilon < \varepsilon_0 \tag{1.5}$$

implies

$$\mathrm{Res}(P) \cap D(\lambda_0, \varepsilon^\gamma) \neq \emptyset. \tag{1.6}$$

Example. Suppose that $P(t)$ is the family of operators defined by choosing $\ell_j = \ell_j(t) \in C^1(\mathbb{R})$, and that $\lambda_0 > 0$ is an eigenvalue of $P(0)$. Then for any $\gamma < 1$ there exists t_0 such that for $|t| \leq t_0$

$$\mathrm{Res}(P(t)) \cap D(\lambda_0, t^\gamma) \neq \emptyset. \tag{1.7}$$

Proof. Let u^0 be a normalized eigenfunction of $P(0)$ with eigenvalue λ_0; in particular $u_k^0 \equiv 0$, $1 \leq k \leq K$. Choose $\chi_j \in C^\infty(\mathbb{R}; [0, 1])$, $j = 1, 2$, such that $\chi_0 + \chi_1 = 1$, $\chi_j(s) = 1$ near $|j - s| < \frac{1}{3}$ and define $u_m^-(x) := \chi_0(x/\ell_m)u_m^0(x)$ and $u_m^+(x) := \chi_1(x/\ell_m)u_m^0(x)$, $u^0 = u^+ + u^-$.

We now define a quasimode for $P(t)$, $u = u(t)$ needed in (1.5):

$$u_m(t) = u_m^-(x) + u_m^+(x - \delta_m(t)), \quad \delta_m(t) := \ell_m(t) - \ell_m(0).$$

For t small enough $\operatorname{supp} u_m^- \subset [0,\frac{2}{3}) \subset [0,\ell_m(t))$ and $\operatorname{supp} u_m^+ \subset (\frac{2}{3},\ell_m(0)] \subset (|\delta_m(t)|,\ell_m(0)]$. Hence the values of $u_m(t)$ and $\partial_\nu u_m(t)$ at the vertices are the same as those of u_m^0 and $u_m(t) \in \mathcal{D}(P(t))$. Also, since $(-\partial_x^2 - \lambda_0^2)u_m^0 = 0$ and $\chi_0^{(k)} = -\chi_1^{(k)}$, (and putting $\ell_m = \ell_m(0)$)

$$\begin{aligned}[(P(t)-\lambda_0^2)u(t)]_m = \ell_m^{-2}(\chi_0''((x-\delta_m(t))/\ell_m)u_m^0(x-\delta_m(t)) - \chi_0''(x/\ell_m)u_m^0(x))\\ + 2\ell_m^{-1}(\chi_0'((x-\delta_m(t))/\ell_m)u_m^0(x-\delta_m(t)) - \chi_0'(x/\ell_m)u_m^0(x)).\end{aligned}$$

We note that all the terms are supported in $(\frac{1}{3}-|\delta_m(t)|,\frac{2}{3}+|\delta_m(t)|)$ and elementary estimates show that $\|(P(t)-\lambda_0^2)u(t)\| \le Ct$. For instance,

$$\begin{aligned}\|\chi_0''(x)(u_m^0(x-\delta_m(t)) - u_m^0(x))\| &\le C'|\delta_m(t)| \max_{|x-\frac{1}{2}|\le \frac{1}{6}+|\delta_m(t)} |\partial_x u_m^0(x)|\\ &\le C'|\delta_m(t)|(\|-\partial_x^2 u_0^m\|_{L^2((\frac{1}{4},\frac{3}{4}))} + \|u_0^m\|_{L^2((\frac{1}{4},\frac{3}{4}))})\\ &\le C''(\lambda_0^2+1)t.\end{aligned}$$

From (1.6) we conclude (after decreasing γ and t_0) that (1.7) holds. □

Remarks. 1. A slightly sharper statement than (1.6) can already be obtained from the proof in §4. It is possible that in fact $\operatorname{Res}(P) \cap D(\lambda_0, C_0\varepsilon)$ where C_0 depends on $\mathcal{L}, R$ and δ. That is suggested by the fact that the converse to this stronger conclusion is valid – see Proposition 4.5. This improvement would require finer complex analytic arguments. It is interesting to ask if methods more specific to quantum graphs, in place of our general methods, could produce this improvement.

2. By adapting Stefanov's methods [15] one can strengthen the conclusion by adding adding a statement about multiplicities (see also [8, Exercise 7.1]) but again we opted for a simple presentation.

Acknowledgements. We are grateful for the support of National Science Foundation under the grant DMS-1500852. We would also like to thank Semyon Dyatlov for helpful discussions and assistance with figures.

2. A Fermi golden rule for boundary value problems: surfaces with cusps

To illustrate the Fermi golden rule in the setting of boundary value problems we consider surfaces, X, with cusps of constant negative curvature. That means that (X,g) is a surface with a smooth boundary and a decomposition (see Fig. 3)

$$\begin{gathered}X = X_1 \cup X_0, \quad \partial X_0 = \partial X_1 \cup \partial X, \quad \partial X_1 \cap \partial X_0 = \emptyset,\\ (X_1, g|_{X_1}) \simeq ([a,\infty)_r \times (\mathbb{R}/\ell\mathbb{Z})_\theta, dr^2 + e^{-2r}d\theta^2).\end{gathered} \tag{2.1}$$

We consider the following family of unbounded operators on $L^2(X)$:

$$P(t) = -\Delta_g - \tfrac{1}{4}, \quad \mathcal{D}(P(t)) = \{u \in H^2(X) : \partial_\nu u|_{\partial X} = \gamma(t)u|_{\partial X}\}. \tag{2.2}$$



FIGURE 3. A surface with one cusp end and a boundary. Suppose we consider a family of boundary conditions for the Laplacian $-\Delta$: $\partial_\nu w = \gamma(t) w$ at ∂X. The Laplacian has continuous spectrum with a family of generalized eigenfuctions $e(\lambda) \in C^\infty(X)$ – see (2.3). Suppose that for $t = 0$, λ^2 is a *simple embedded* eigenvalue of $-\Delta$ with the boundary condition $\partial_\nu w = \gamma(0) w$, with the normalized eigenfunction given by u. Then $\lambda = \lambda(0)$ belong to a smooth family of *resonances* of Laplacians with boundary condition $\partial_\nu w = \gamma(t) w$, and $\operatorname{Im} \ddot\lambda = -\frac{1}{4\lambda^2} |\langle \dot\gamma u, e(\lambda)\rangle|^2$ – see Theorem 3.

where $t \mapsto \gamma(t) \in C^\infty(\partial X)$ is a smooth family of functions on ∂X and ∂_ν is the outward pointing normal derivative. The spectrum of the operator P has the following well known decomposition:

$$\begin{gathered} \operatorname{Spec}(P) = \operatorname{Spec}_{\rm pp}(P) \cup \operatorname{Spec}_{\rm ac}(P), \quad \operatorname{Spec}_{\rm ac}(P) = [0, \infty), \\ \operatorname{Spec}_{\rm pp}(P) = \{E_j\}_{j=0}^J, \quad -\tfrac14 \le E_0 < E_1 \le E_2 \cdots, \quad 0 \le J \le +\infty. \end{gathered}$$

(When $J = +\infty$ then $E_j \to \infty$.) The eigenvalues $E_j > 0$ are *embedded* in the continuous spectrum. In addition the resolvent $R(\lambda) := (P - \lambda^2)^{-1} : L^2 \to L^2$, $\operatorname{Im} \lambda > 0$, has a meromorphic continuation to $\lambda \in \mathbb{C}$ as an operator $R(\lambda) : C_{\rm c}^\infty(X) \to C^\infty(X)$. Its poles are called *scattering resonances*. Under generic perturbation of the metric in X_0 all embedded eigenvalues become resonances. For proofs of these well known facts see [4] and also [8, §4.1 (Example 3), §4.2 (Example 3), §4.4.2] for a presentation from the point of view of *black box scattering* [13].

The generalized eigenfunctions, $e(\lambda, x)$, describing the projection onto the continuous spectrum have the following properties:

$$\begin{gathered} (P - \lambda^2) e(\lambda, x) = 0, \qquad \frac{1}{\ell} \int_0^\ell e(\lambda, x)|_{X_1} d\theta = e^{\frac{r}{2}} \left(e^{-i\lambda r} + s(\lambda) e^{i\lambda r} \right), \\ (R(\lambda) - R(-\lambda)) f = \tfrac{i}{2\lambda} e(\lambda, x) \langle f, e(\lambda, \bullet)\rangle, \quad \lambda \in \mathbb{R}, \quad f \in C_{\rm c}^\infty(X), \end{gathered} \tag{2.3}$$

see [8, Theorem 4.20]. With these preliminaries in place we can now prove

Theorem 3. *Suppose that the operators $P(t)$ are defined by (3.3) and that $\lambda > 0$ is a* simple *eigenvalue of $P(0)$ and $(P(0) - \lambda^2) u = 0$, $\|u\|_{L^2} = 1$.*

Then there exists a smooth function $t \mapsto \lambda(t)$, $|t| < t_0$, such that $\lambda(0) = \lambda$, $\lambda(t)$ is a scattering resonance of $P(t)$ and

$$\operatorname{Im} \ddot\lambda = -\frac{1}{4\lambda^2} \left| \langle \dot\gamma u, e\rangle_{L^2(\partial X)} \right|^2, \quad e(x) = e(\lambda, x), \tag{2.4}$$

where $e(\lambda, x)$ is given in (2.3), $\dot f := \partial_t f|_{t=0}$ and $L^2(\partial X)$ is defined using the metric induced by g.

Remarks. 1. For recent advances in mathematical study of the Fermi golden rule in more standard settings of mathematical physics and for numerous references see Cornean–Jensen–Nenciu [5].

2. In the case of scattering on constant curvature surfaces with cusps the Fermi golden rule was explicitly stated by Phillips–Sarnak – see[12] and for a recent discussion [11]. For a presentation from the black box point of view see [8, §4.4.2].

3. The proof generalizes immediately to the case of several cusps (which is analogous to a quantum graph with several leads), $(X_k, g|_{X_k}) \simeq ([a_k, \infty) \times \mathbb{R}/\ell_k\mathbb{Z}, dr^2 + e^{-2r} d\theta^2$, $1 \le k \le K$. In that case the generalized eigenfunction are normalized using

$$\frac{1}{\ell_m} \int_0^{\ell_m} e^k(\lambda, x)|_{X_m} d\theta = e^{\frac{r}{2}} \left(\delta_{km} e^{-i\lambda r} + s_{km}(\lambda) e^{i\lambda r} \right).$$

The Fermi golden rule for the boundary value problem (3.3) is given by

$$\operatorname{Im} \ddot\lambda = -\frac{1}{4\lambda^2} \sum_{k=1}^{K} \left| \langle \dot\gamma u, e^k\rangle_{L^2(\partial X)} \right|^2, \quad e^k(x) = e^k(\lambda, x). \tag{2.5}$$

Proof. For notational simplicity we assume that $\gamma(0) \equiv 0$, that is that $P(0)$ is the Neumann Laplacian on X. We will also omit the parameter t when that is not likely to cause confusion. It is also convenient to use $z = \lambda^2$ and to write $\langle \bullet, \bullet \rangle$ for the $L^2(X, d\operatorname{vol}_g)$ inner product and $\langle \bullet, \bullet \rangle_{L^2(\partial X)}$ for the inner product on $L^2(\partial)$ with the measure induced by the metric g.

We first define the following orthogonal projection:

$$\begin{gathered} \mathbb{1}_{r\ge R}\, u := \frac{1}{\ell} \int_0^{\ell} u|_{X_1 \cap \{r \ge R\}}\, d\theta, \qquad \mathbb{1}_{r\ge R} : L^2(X) \to L^2([R, \infty), e^{-r} dr), \quad R > a, \\ \mathbb{1}_{r\le R} := I - \mathbb{1}_{r\ge R}, \quad \mathcal{H}_R := \mathbb{1}_{r\le R}\, L^2(X). \end{gathered} \tag{2.6}$$

The smoothness of scattering resonances arising from a smooth perturbation of a simple resonance follows from smooth dependence of the continuation of $(P(t) - \lambda^2)^{-1}$ (see Proposition 4.3 below for a general argument). Let $t \mapsto u(t)$, $u(0) = u$ denote a smooth

family of resonant states:

$$\begin{gathered}(P(t)-z(t))u(t)=0,\quad \frac{1}{\ell}\int_0^\ell u(t)|_{X_1}d\theta=a(t)e^{\frac{r}{2}}e^{i\lambda(t)r},\\ a(0)=0,\quad \operatorname{Im}\lambda(t)\leq 0,\quad \lambda(0)^2=z(0).\end{gathered}\tag{2.7}$$

The second equation in (2.7) means that $u(t)$ is *outgoing* – see [8, §4.4].

The self-adjointness of $P(t)$ and integration by parts for the zero mode in the cusp show that for $u=u(t)$ and $P=P(t)$,

$$\begin{aligned}0&=\operatorname{Im}\langle (P-z)u,\mathbb{1}_{r\leq R}\,u\rangle\\ &=-\operatorname{Im}\partial_r(\mathbb{1}_{r\geq R}\,u)(R)\overline{\mathbb{1}_{r\geq R}\,u}(R)-\operatorname{Im}z\|\,\mathbb{1}_{r\leq R}\,u\|^2_{L^2(X)}.\end{aligned}\tag{2.8}$$

(See [8, (4.4.17)] for a detailed presentation in the general black box setting.) Since $\operatorname{Im}\dot z=0$ (as $\operatorname{Im}z(t)\leq 0$, see also (2.12) below) and since $\mathbb{1}_{r\geq R}\,u(0)=0$, we have have, at $t=0$, $\operatorname{Im}\ddot z=-2\operatorname{Im}\partial_r(\mathbb{1}_{r\geq R}\,\dot u)(R)\overline{\mathbb{1}_{r\geq R}\,\dot u}$. We would like to argue as in (2.8) but in reverse. However, as $\dot u$ will not typically be in $\mathcal{D}(P)$ we now obtain boundary terms:

$$\operatorname{Im}\ddot z=2\operatorname{Im}\langle (P-z)\dot u,\mathbb{1}_{r\leq R}\,\dot u\rangle+2\operatorname{Im}\langle\partial_\nu\dot u,\dot u\rangle_{L^2(\partial X)}.\tag{2.9}$$

We now need an expression for $\dot u$. Since $(P(t)-z(t))u(t)=0$, $\partial_\nu u|_{\partial X}=\gamma u|_{\partial X}$, we have (at $t=0$),

$$(P-z)\dot u=\dot z u,\quad \partial_\nu\dot u|_{\partial X}=\dot\gamma u|_{\partial X}.\tag{2.10}$$

In addition, differentiation of the second condition in (2.7) shows that $\dot u$ is outgoing.

Without loss of generality we can assume that $u=u(0)$ is real valued. Choose $g\in\bar C^\infty(X,\mathbb{R})$ (real valued, compactly supported and smooth up to the boundary) such that $\partial_\nu g|_{\partial X}=\dot\gamma u|_{\partial X}$. We claim that

$$\langle \dot z u-(P-z)g,u\rangle=0.\tag{2.11}$$

In fact, Green's formula shows that the left hand side of (2.11) is equal to $\dot z+\int_{\partial X}\dot\gamma u^2$. On the other hand, using the fact that $\mathbb{1}_{r\leq R}\,u(0)=u(0)$,

$$\begin{aligned}0&=-\frac{d}{dt}\langle (P(t)-z(t))u(t),\mathbb{1}_{r\leq R}\,u(t)\rangle|_{t=0}=\langle\dot z u-(P-z)\dot u,u\rangle\\ &=\dot z+\int_{\partial X}\partial_\nu\dot u u=\dot z+\int_{\partial X}\gamma u^2.\end{aligned}\tag{2.12}$$

In view of (2.11), $v:=g+R(\lambda)(\dot z-(P-z)g)$, $\lambda^2=z$, $\lambda>0$, is well defined, outgoing (see (2.7)) and solves the boundary value problem (2.10) satisfied by $\dot u$. Since the eigenvalue at z is simple that means that $\dot u-v$ is a multiple of u (see [8, Theorem 4.18] though in this one dimensional case this is particularly simple). Hence

$$\dot u=\alpha u+g+R(\lambda)(\dot z u-(P-z)g).\tag{2.13}$$

With this formula in place we return to (2.9). First we note that the first term on the right hand side vanishes:

$$\begin{aligned}\operatorname{Im}\langle (P-z)\dot{u}, \mathbb{1}_{r\leq R}\,\dot{u}\rangle &= \operatorname{Im}\langle \dot{z}u, \dot{u}\rangle = \dot{z}\operatorname{Im}\langle u, \alpha u + g + R(\lambda)(\dot{z}u - (P-z)g)\rangle \\ &= \dot{z}\operatorname{Im}\alpha + \dot{z}\operatorname{Im}\langle u, R(\lambda)(\dot{z}u-(P-z)g)\rangle \\ &= \dot{z}\operatorname{Im}\alpha.\end{aligned} \tag{2.14}$$

Here we used the fact that u and g were chosen to be real. The last identity followed from (2.11). To analyse the second term on the right hand side of (2.9) we recall some properties of the Schwartz kernel of the resolvent:

$$R(\lambda)(x,y) = R(\lambda)(y,x) = \overline{R(-\overline{\lambda})(x,y)}, \quad \lambda\in\mathbb{C}. \tag{2.15}$$

(The first property follows from considering $\lambda = ik$, $k\gg 1$, and using the fact that $\overline{Pu} = P\bar{u}$, and the second from considering $\operatorname{Im}\lambda \gg 1$, $z=\lambda^2$, and noting that $((P-z)^{-1})^* = (P-\bar{z})^{-1}$.) Using (2.9),(2.10),(2.14),(2.13),(2.15),(2.12) and the fact that u and g are real, we now see that

$$\begin{aligned}\operatorname{Im}\ddot{z} &= 2\dot{z}\operatorname{Im}\alpha + 2\operatorname{Im}\langle \dot{\gamma}u, \dot{u}\rangle_{L^2(\partial X)} \\ &= 2\dot{z}\operatorname{Im}\alpha + 2\operatorname{Im}\alpha\langle\dot{\gamma}u, u\rangle + 2\operatorname{Im}\langle\dot{\gamma}u, [R(\lambda)(\dot{z}u-(P-z)g)]|_{\partial X}\rangle_{L^2(\partial X)} \\ &= \tfrac{1}{i}\langle\dot{\gamma}u, [(R(\lambda)-R(-\lambda))(\dot{z}u-(P-z)g)]|_{\partial X}\rangle_{L^2(\partial X)}.\end{aligned} \tag{2.16}$$

Since $(R(\lambda)-R(-\lambda))u = 0$ we have now use (2.3) to see that

$$\begin{aligned}[(R(\lambda)-R(-\lambda))(\dot{z}u-(P-z)g)]|_{\partial X} &= -\tfrac{i}{2\lambda}e(\lambda)|_{\partial X}\int_X \overline{e(\lambda)}(P-z)g \\ &= -\tfrac{i}{2\lambda}e(\lambda)|_{\partial X}\int_{\partial X}(\partial_\nu\bar{e}(\lambda)g - \partial_\nu g\overline{e(\lambda)}) \\ &= \tfrac{i}{2\lambda}e(\lambda)|_{\partial X}\langle\dot{\gamma}u, e\rangle_{L^2(\partial X)}.\end{aligned}$$

Inserting this into (2.16) gives (2.4) completing the proof. □

3. Proof of Theorem 1

We follow the same strategy as in the proof of Theorem 3 but with some notational complexity due to the graph structure.

Let $H^2 := \bigoplus_{m=1}^{M+K} H^2([0,\ell_m])$. Then for $u,v\in H^2$, $(\partial_x^k u)_m := \partial_x^k u_m$,

$$\begin{aligned}-\langle\partial_x^2 f, g\rangle_{L^2} &= \langle\partial_x f, \partial_x g\rangle_{L^2} - \sum_v\sum_{e_m\ni v}\partial_\nu f_m(v)\bar{g}_m(v) \\ &= -\langle f, \partial_x^2 g\rangle_{L^2} + \sum_v\sum_{e_m\ni v}\big(f_m(v)\partial_\nu\bar{g}_m(v) - \partial_\nu f_m(v)\bar{g}_m(v)\big).\end{aligned} \tag{3.1}$$

We note here that the sum over vertices can be written as a sum over edges:

$$\sum_v \sum_{e_m \ni v} \partial_\nu f_m(v)\bar{g}_m(v) = \sum_{m=1}^{M+K} \sum_{v\in\partial e_m} \partial_\nu f_m(v)\bar{g}_m(v). \tag{3.2}$$

Just as in §2 the domain of the deformed operators will change but we make a modification which will keep the Hilbert space on which $\widetilde{P}(t)$ (we change the notation from §1 and will use $P(t)$ for a unitarily equivalent operator) acts fixed by changing the lengths in (1.2). For that let

$$L^2_t := \bigoplus_{m=1}^{M+K} L^2([0, e^{-a_m(t)}\ell_m]), \quad L^2 := L^2_0, \quad U(t) : L^2_t \to L^2,$$
$$[U(t)u]_m(y) := e^{-a_m(t)/2} u_m(e^{-a_m(t)}y), \quad U(t)^{-1} = U(t)^*.$$

Let $\widetilde{P}(t)$ be defined in L^2_t by $(\widetilde{P}(t)u)_m = -\partial_x^2 u_m$,

$$\mathcal{D}(\widetilde{P}(t)) = \{u : u_m \in H^2([0, e^{-a_j(t)}\ell_m]),\ u_m(v) = u_\ell(v),\ v \in e_m \cap e_\ell,\ \sum_{e_m\ni v} \partial_\nu u_m(v) = 0\}.$$

That is just the family of Neumann Laplace operators on the graph with the lengths $e^{-a_j(t)}\ell_j$.

On L^2 we define a new family of operators: $P(t) := U(t)\widetilde{P}(t)U(t)^*$. It is explicitly given by $[P(t)u]_m = -e^{2a_m(t)}\partial_x^2 u_m$,

$$\begin{aligned}\mathcal{D}(P(t)) = \{u \in H^2 : e^{a_m(t)/2}u_m(v) = e^{a_\ell(t)/2}u_\ell(v),\ v \in e_m \cap e_\ell,\\ \sum_{e_m\ni v} e^{3a_m(t)/2}\partial_\nu u_m(v) = 0\}.\end{aligned} \tag{3.3}$$

Using Proposition 4.3 from the next section we see that for small t there exists a smooth family $t \mapsto u(t) \in H^2_{\rm loc}$ such that

$$\begin{gathered}(P(t) - z(t))u(t) = 0, \quad u_k(t,x) = a(t)e^{i\lambda(t)x}, \quad k \geq M+1,\\ \operatorname{Im}\lambda(t) \leq 0, \quad \lambda(0)^2 = z, \quad \lambda(0) > 0.\end{gathered} \tag{3.4}$$

We defined $\mathcal{H}_R$ by (1.4) and denote by $\mathbb{1}_{x\leq R}$ the orthogonal projection $L^2 \to \mathcal{H}_R$.

Writing $P = P(t)$, $u = u(t)$, $z = z(t)$ we see, as in (2.8), that

$$0 = \operatorname{Im}\langle (P-z)u, \mathbb{1}_{x\leq R}\, u\rangle = -\operatorname{Im}\sum_{m=1}^K \partial_x u_m(R)\bar{u}_m(R) - \operatorname{Im} z\|u\|^2_{\mathcal{H}_R}. \tag{3.5}$$

We recall that e_m, $1 \le m \le K$ are the infinite edges with unique boundaries. Hence, using (3.1), at $t = 0$,

$$\begin{aligned} \operatorname{Im} \ddot{z} &= 2 \operatorname{Im} \sum_{m=1}^{K} \partial_x \dot{u}_m(R) \overline{\dot{u}}_m(R) \\ &= 2 \operatorname{Im} \langle (P - z)\dot{u}, \mathbb{1}_{x \le R}\, \dot{u} \rangle + 2 \operatorname{Im} \sum_v \sum_{e_m \ni v} \partial_\nu \dot{u}_m(v) \overline{\dot{u}}_m(v). \end{aligned} \tag{3.6}$$

We now look at the equation satisfied by $\dot{u}$ at $t = 0$:

$$\frac{d}{dt}(P(t) - z(t))u(t) = 2\dot{a}(-\partial_x^2 u) - \dot{z}u + (P - z)\dot{u} = (2\dot{a}z - \dot{z})u + (P - z)\dot{u}. \tag{3.7}$$

Hence,

$$\begin{gathered} (-\partial_x^2 - z)\dot{u}_m = (\dot{z} + 2z\dot{a}_m)u_m, \quad \sum_{e_m \ni v} \partial_\nu \dot{u}_m(v) = -\tfrac{3}{2} \sum_{e_m \ni v} \dot{a}_m \partial_\nu u_m(v), \\ \dot{u}_m(v) - \dot{u}_\ell(v) = \tfrac{1}{2}(\dot{a}_\ell - \dot{a}_m)u(v), \quad v \in e_m \cap e_\ell. \end{gathered} \tag{3.8}$$

We used here the fact that $u(v) := u_m(v)$ does not depend on m. The second condition can be formulated as $\dot{u}_m(v) = w(v) - \frac{1}{2}\dot{a}_m(v)u(v)$, where $w := \partial_t(e^{a(t)/2}u(t))|_{t=0}$ is continuous on the graph.

To find an expression for $\dot{u}$ (similar to (2.13)) we first find

$$g \in \bigoplus_{m=1}^{K} C_c^\infty([0, \infty)) \oplus \bigoplus_{m=K+1}^{M+K} C^\infty([0, \ell_m]),$$

such that

$$\sum_{e_m \ni v} \partial_\nu g_m(v) = -\tfrac{3}{2} \sum_{e_m \ni v} \dot{a}_m u_m(v), \quad g_m(v) - g_\ell(v) = \tfrac{1}{2}(\dot{a}_\ell - \dot{a}_m)u(v). \tag{3.9}$$

We can assume without loss of generality that both g and u are real valued.

In analogy to (2.11) we claim that

$$\langle (\dot{z} - 2z\dot{a})u - (P - z)g, u \rangle = 0. \tag{3.10}$$

In fact, using (3.1), (3.7) and (3.8) we obtain

$$\begin{aligned} 0 &= -\frac{d}{dt}\langle (P(t) - z(t))u(t), \mathbb{1}_{x \le R}\, u(t) \rangle|_{t=0} = \langle \dot{z}u - 2z\dot{a}u - (P - z)\dot{u}, u \rangle \\ &= \dot{z} - 2z\langle \dot{a}u, u \rangle + \sum_v \sum_{e_m \ni v} (\partial_\nu \dot{u}_m(v)u(v) - \dot{u}_m(v)\partial_\nu u_m(v)) \\ &= \dot{z} - 2z\langle \dot{a}u, u \rangle + \sum_v \sum_{e_m \ni v} (-\tfrac{3}{2}\dot{a}_m \partial_\nu u_m(v)u(v) - (w(v) - \tfrac{1}{2}\dot{a}_m u(v))\partial_\nu u_m(v)) \\ &= \dot{z} - 2z\langle \dot{a}u, u \rangle - \sum_v \sum_{e_m \ni v} \dot{a}_m \partial_\nu u_m(v)u(v). \end{aligned} \tag{3.11}$$

(We used the continuity of u and the Neumann condition $\sum_{e_m\ni v}\partial_\nu u_m(v)=0$.) Since g and u satisfy the same boundary conditions (3.8) and (3.9) (3.10) follows from (3.11).

As in the derivation of (2.13) we now see that for some $\alpha\in\mathbb{C}$ we have

$$\dot u=\alpha u+g+R(\lambda)(\dot z u-2z\dot a u-(P-z)g). \tag{3.12}$$

With this in place we return to (3.6). The first term on the right hand side is

$$\begin{aligned}2\operatorname{Im}\langle(P-z)\dot u,\mathbb{1}_{x\leq R}\,\dot u\rangle&=2\operatorname{Im}\langle\dot z u-2z\dot a u,\dot u\rangle\\&=2\operatorname{Im}\langle\dot z u-2z\dot a u,\alpha u+g+R(\lambda)(\dot z u-2z\dot a u-(P-z)g)\rangle\\&=2\operatorname{Im}\alpha(\dot z-2z\langle\dot a u,u\rangle)\\&\qquad-4z\operatorname{Im}\langle\dot a u,R(\lambda)(\dot z u-2z\dot a u-(P-z)g)\rangle.\end{aligned}$$

(We used here the simplifying assumption that g and u are real valued.)

As in (2.16) we conclude that

$$4z\operatorname{Im}\langle\dot a u,R(\lambda)(-\dot z u+2z\dot a u+(P-z)g)\rangle=\tfrac{2z}{i}\langle\dot a u,[(R(\lambda)-R(-\lambda)](2z\dot a u+(P-z)g)\rangle$$

Now, as in (2.3), [8, Theorem 4.20] shows that

$$(R(\lambda)-R(-\lambda))f=\frac{i}{2\lambda}\sum_{k=1}^K e^k(\lambda,x)\langle f,e^k(\lambda,\bullet)\rangle,\quad \lambda\in\mathbb{R},\quad f\in\mathcal{H}_R, \tag{3.13}$$

which means that (with $z=\lambda^2$ and $e^k=e^k(\lambda)$)

$$\begin{aligned}&\tfrac{2z}{i}\langle\dot a u,[(R(\lambda)-R(-\lambda)](2z\dot a u+(P-z)g)\rangle=\\&\qquad-2\lambda^3\sum_{k=1}^K|\langle\dot a u,e^k\rangle|^2-\lambda\sum_{k=1}^K\langle\dot a u,e^k\rangle\langle e^k(\lambda),(P-z)g\rangle.\end{aligned}$$

The second term on the right hand side is now rewritten using (3.1) and the boundary conditions (3.9):

$$\begin{aligned}&\lambda\sum_{k=1}^K\langle\dot a u,e^k\rangle\left(\sum_v\sum_{e_m\ni v}(\partial_\nu e^k_m(v)g_m(v)-\partial_\nu g_m(v)e^k(v))\right)=\\&\qquad\lambda\sum_{k=1}^K\langle\dot a u,e^k\rangle\left(\sum_v\sum_{e_m\ni v}\tfrac12\dot a_m(-\partial_\nu e^k_m(v)u(v)+3\partial_\nu u_m(v)e^k(v))\right).\end{aligned}$$

We conclude that

$$\begin{aligned}2\operatorname{Im}\langle(P-z)\dot u,\mathbb{1}_{x\leq R}\,\dot u\rangle&=2\operatorname{Im}\alpha(\dot z-2z\langle\dot a u,u\rangle)-2\lambda^3\sum_{k=1}^K|\langle\dot a u,e^k\rangle|^2\\&\quad-2\lambda\sum_{k=1}^K\langle\dot a u,e^k\rangle\left(\sum_v\sum_{e_m\ni v}\tfrac14\dot a_m(3\partial_\nu u_m(v)e^k(v)-\partial_\nu e^k_m(v)u(v))\right).\end{aligned} \tag{3.14}$$

A similar analysis of the second term on the right hand side of (2.9) shows that

$$2\,\mathrm{Im}\sum_v\sum_{e_m\ni v}\partial_\nu\dot u_m(v)\overline{\dot u}_m(v)=\mathrm{Im}\,\alpha\left(2\sum_v\sum_{e_m\ni v}\dot a_m\partial_\nu u_m(v)u(v)\right)$$
$$-2\lambda^{-1}\sum_{k=1}^K\left|\sum_v\sum_{e_m\ni v}\tfrac14\dot a_m(\partial_\nu e_m^k(v)u(v)-3\partial_\nu u_m(v)e^k(v))\right|^2 \tag{3.15}$$
$$-2\lambda\sum_{k=1}^K\overline{\langle\dot a u,e^k\rangle}\left(\sum_v\sum_{e_m\ni v}\tfrac14\dot a_m(3\partial_\nu u_m(v)\overline{e}^k(v)-\partial_\nu\overline{e}_m^k(v)u(v))\right).$$

Inserting (3.14),(3.15) into (3.6), using (3.11) and $\mathrm{Im}\,\ddot z=2\lambda\,\mathrm{Im}\,\ddot\lambda$ gives (1.3). □

Example 1. Consider a connected graph with M bonds and K leads. Suppose that an embedded eigenvalue λ is simple and satisfies

$$\lambda\ell_m\in\pi\mathbb{Z},\ m=1,\cdots,M. \tag{3.16}$$

Then

$$\mathrm{Im}\,\ddot\lambda=-\sum_{k=1}^K\left|\lambda\langle\dot a u,e^k(\lambda)\rangle\right|^2. \tag{3.17}$$

Proof. $u_m(x)=C_m\sin(\lambda x)$ where e_m and a lead are meeting at a vertex. Since the graph is connected, $u_m(x)=C_m\sin(\lambda x)$ for $1\le m\le M$. Let $n_m=\frac{\lambda\ell_m}{\pi}$ and let

$$e_m^k(\lambda,x)=A_{mk}\sin(\lambda x)+B_{mk}\cos(\lambda x).$$

Then $u_m(\ell_m)=u_m(\ell_m)=0$ and

$$\partial_\nu u_m(0)=(-1)^{n_m+1}\partial_\nu u_m(\ell_m),\ e_m^k(\lambda,\ell_m)=(-1)^{n_m}e_m^k(\lambda,0).$$

We can use this and (3.2) to reduce F_k in (1.3) to

$$F_k=\lambda\langle\dot a u,e^k(\lambda)\rangle+\lambda^{-1}\sum_{m=1}^M\tfrac34\dot a_m(\partial_\nu u_m(0)\overline{e^k(\lambda,0)}+\partial_\nu u_m(\ell_m)\overline{e^k(\lambda,\ell_m)})$$
$$=\lambda\langle\dot a u,e^k(\lambda)\rangle.$$

Theorem 1 then gives (3.17). □

Example 2. Let us consider a graph with $M=5,K=2$ and four vertices: see in Fig. 4. Let $\ell_m(0)=1,\ 1\le m\le 5$. Then the sequence of embedded eigenvalues λ is given as $S_1\cup S_2$ where

$$S_1=\pi\mathbb{Z},\quad S_2=\left\{\lambda\ :\ \tan\lambda+2\tan\tfrac{\lambda}{2}=0,\quad \lambda\notin\tfrac{\pi}{2}\mathbb{Z}\right\}.$$

If $\lambda\in S_1$, then (3.16) is satisfied. If $\lambda\in S_2$, however, we have (with v_1 and v_2 corresponding to $x=0$ for e_3, e_6 and e_4, e_5 respectively, and v_4 to $x=0$ for e_7)

$$u_3(x)=C\sin(\lambda x),\quad u_4(x)=C\sin(\lambda x),\quad u_5(x)=-C\sin(\lambda x),$$



FIGURE 4. The graph from Example 2: in this case boundary terms in our Fermi golden rule appear at some embedded eigenvalues such as λ_0 which is the smallest solution of $\tan\lambda + 2\tan\frac{\lambda}{2} = 0$, $\lambda_0 \approx 1.9106$. We consider the following variation of length: $\ell_3 = 1-t, \ell_4 = 1+t, \ell_5 = 1-t, \ell_6 = 1+t$, and (a): $\ell_7 = 1$ (b): $\ell_7 = 1 + t/2$ (c): $\ell_7 = 1 + t$.

$$u_6(x) = -C\sin(\lambda x), \quad u_7(x) = C\frac{\sin\lambda}{\sin\frac{\lambda}{2}}\sin\left(\lambda\left(x-\tfrac{1}{2}\right)\right),$$

where $C > 0$ is the normalization constant. Note that

$$u(v_3) = C\sin\lambda \neq 0, \quad u(v_4) = -C\sin\lambda \neq 0.$$

So we do not have the simple formula (3.17) in this case.

4. Proof of Theorem 2

The proof adapts to the setting of quantum graphs and of quasimodes u satisfying (1.5) the arguments of [18]. They have origins in the classical work of Carleman [3] on completeness of eigenfunctions for classes non-self-adjoint operators, see also [15] and [17].

We start with general results which are a version of the arguments of [8, §7.2]. In particular they apply without modification to quantum graphs with general Hamiltonians and general boundary conditions. We note that for metric graphs considered here much more precise estimates are obtained by Davies–Pushnitski [6] and Davies–Exner–Lipovský [7] but since we want uniformity we present an argument illustrating the black box point of view.

Proposition 4.1. *Suppose that P satisfies the assumptions of Theorem 2 and $\Omega_1 \Subset \Omega_2 \Subset \mathbb{C}$, where Ω_j are open sets.*

Then there exist constants C_1 depending only on Ω_2 and $\mathcal{L}$, and C_2 depending on Ω_1, Ω_2, R and $\mathcal{L}$ such that

$$|\operatorname{Res}(P) \cap \Omega_2| \le C_1,$$
$$\| \mathbb{1}_{r\le R}\, R(\lambda)\, \mathbb{1}_{r\le R} \|_{L^2\to L^2} \le C_2 \prod_{\zeta\in \operatorname{Res}(P)\cap\Omega_2} |\lambda-\zeta|^{-1}, \quad \lambda\in\Omega_1, \tag{4.1}$$

where the elements of $\operatorname{Res}(P)$ *are included according to their multiplicities.*

Proof. Let $R_0(\lambda) : \bigoplus_{k=1}^K L^2_{\rm comp}(e_k) \to \bigoplus H^2_{\rm loc}\cap H^1_{0,\rm loc}(e_k)$, be defined as the diagonal operator acting on each component as $R_0^0(\lambda)$, the Dirichlet resolvent on $L^2_{\rm comp}([0,\infty))$ continued *analytically* to all of $\mathbb{C}$:

$$R_0^0(\lambda)f(x) = \int_0^\infty \frac{e^{i\lambda(x+y)} - e^{i\lambda|x-y|}}{2i\lambda} f(y)dy.$$

To describe $\mathbb{1}_{r\le R}\, R(\lambda)\, \mathbb{1}_{r\le R}$ we follow the general argument of [13] (see also [8, §4.2,4.3]). For that we choose $\chi_j \in C^\infty_{\rm c}$, $j = 0, \cdots, 3$ to be equal to 1 on all edges and to satisfy

$$\chi_j|_{e_k} \in C^\infty_{\rm c}([0,2R)), \quad \chi_0|_{e_k}(x) = 1,\ x\le R, \quad \chi_j|_{e_k}(x) = 1,\ x\in \operatorname{supp}\chi_{j-1}|_{e_k},$$

for $k = 1, \cdots, K$. For λ_0 with $\operatorname{Im}\lambda_0 > 0$, we define

$$Q(\lambda,\lambda_0) := (1-\chi_0)R_0(\lambda)(1-\chi_1) + \chi_2 R(\lambda_0)\chi_1, \quad Q(\lambda,\lambda_0) : L^2_{\rm comp} \to \mathcal{D}_{\rm loc}(P).$$

Then

$$(P-\lambda^2)Q(\lambda,\lambda_0) = I + K(\lambda,\lambda_0),$$
$$K_0(\lambda,\lambda_0) := -[P,\chi_0]R_0(\lambda)(1-\chi_1) + (\lambda_0^2-\lambda^2)\chi_2 R(\lambda_0)\chi_1 + [P,\chi_2]R(\lambda_0)\chi_1.$$

We now choose $\lambda_0 = e^{\pi i/4}\mu$, $\mu \gg 1$. Then

$$I + K_0(\lambda_0,\lambda_0) \ \text{ and } \ I + K_0(\lambda_0,\lambda_0)\chi_3 \ \text{ are invertible on } \ L^2, \tag{4.2}$$

$K(\lambda,\lambda_0)\chi_3$ is compact, and

$$R(\lambda) = Q(\lambda,\lambda_0)(I + K_0(\lambda,\lambda_0)\chi_3)^{-1}(I - K_0(\lambda,\lambda_0)(1-\chi_3)), \tag{4.3}$$

where $\lambda \mapsto (I + K_0(\lambda,\lambda_0)\chi_3)^{-1}$ is a meromorphic family of operators. We now put

$$K(\lambda,\lambda_0) := K_0(\lambda,\lambda_0)\chi_3$$

and conclude that

$$\mathbb{1}_{r\le R}\, R(\lambda)\, \mathbb{1}_{r\le R} = \mathbb{1}_{r\le R}\, Q(\lambda,\lambda_0)\chi_3 (I + K(\lambda,\lambda_0))^{-1}\, \mathbb{1}_{r\le R}, \tag{4.4}$$

and the set of resonances is given by the poles of $(I + K(\lambda,\lambda_0))^{-1}$. (See [8, §4.2] and in particular [8, (4.2.19)].)

We now claim that $K(\lambda,\lambda_0)$ is of trace class for $\lambda\in\mathbb{C}$ and that for a any compact subset $\Omega \Subset \mathbb{C}$ there exists a constant C_3 depending only on Ω, $\mathcal{L}$ and λ_0 such that

$$\|K(\lambda,\lambda_0)\|_{\rm tr} \le C_3. \tag{4.5}$$

To see this, let $\widetilde{P}$ be the operator of $\mathcal{H}_{3R}$ where we put, say the Neumann boundary condition at $3R$ on each infinite lead. Let $\widetilde{P}_{\min}, \widetilde{P}_{\max}$ be the same operators but on metric graphs were all the length $\ell_j \in \mathcal{L}$, $K+1 \leq j \leq K+M$ were replaced by $\ell_{\min} := \min \mathcal{L}$ and $\ell_{\max} := \max \mathcal{L}$ respectively. These operators have discrete spectra and the ordered eigenvalues of these operators satisfy

$$\lambda_p(\widetilde{P}_{\max}) \leq \lambda_p(\widetilde{P}) \leq \lambda_p(\widetilde{P}_{\min}). \tag{4.6}$$

This is a consequence of the following lemma:

Lemma 4.2. *Suppose that the unbounded operator $\widetilde{P}_k(t) : \mathcal{H}_\rho \to \mathcal{H}_\rho$, $\rho > 0$, with edge length given by*

$$\ell_k(t) = \rho, \ 1 \leq k \leq K, \quad \ell_m(t) = e^{-\delta_{mk} t} \ell_m, \quad K+1 \leq m \leq M+K,$$

and

$$\mathcal{D}(\widetilde{P}_k(t)) = \{u : u_m \in H^2([0, \ell_m(t)]), \ u_m(v) = u_\ell(v), \ v \in e_m \cap e_\ell, \ \sum_{e_m \ni v} \partial_\nu u_m(v) = 0\}.$$

If $0 = \mu_0(t) \leq \mu_1(t) \leq \mu_2(t) \cdots$, is the ordered sequence of eigenvalues of $P(t)$, then $\mu_p(t)$ is an increasing function of t.

Proof. From [2, Theorem 3.10] we know that if μ is an eigenvalue of $P(s)$ of multiplicity N then we can choose analytic functions $\mu^n(t) \in \mathbb{R}$, $u^n(t) \in \mathcal{D}(P(s))$, such that $\mu^n(s) = \mu$, and for small $t-s$, $P(t)u^n(t) = \mu^n(t)u^n(t)$, and $\{u^n(t)\}_{n=1}^N$ is an orthonormal setting spanning $\mathbb{1}_{|P(t)-\mu| \leq \varepsilon} L^2$, for $\varepsilon > 0$ small enough. The lemma follows from showing that $\partial_t \mu^n(s) \geq 0$ for any n.

Without loss of generality we can assume that $s = 0$. We can then use the same calculation as in (3.11) with $z = \mu^n(0)$, $a_m(t) = \delta_{km} t$ and $u = u^n(0)$. That gives

$$\mu_p'(0) = 2\mu_p(0)\langle u, u\rangle_{L^2(e_k)} + \sum_{v \in \partial e_k} \partial_\nu u_k(v) u_k(v).$$

Since $u_k(x) = a \sin \sqrt{\mu}_p x + b \cos \sqrt{\mu}_p x$, for some $a, b \in \mathbb{R}$, a calculation shows that

$$\mu_p'(0) = \mu_p(0)\ell_k(a^2 + b^2) \geq 0,$$

completing the proof. □

The inequality (4.6) follows from the lemma as we can change the length of the edges in succession. The Weyl law for $\widetilde{P}$ (see [1]) and the fact that $\widetilde{P}\chi_3 = P\chi_3$ (where χ_3 denotes the multiplication operator), now shows that for any operator $A : L^2 \to \mathcal{D}(P)$,

$$\|\chi_3 A \chi_3\|_{\mathrm{tr}} \leq C_4 \|P\chi_3 A \chi_3\| + C_4 \|\chi_3 A \chi_3\|,$$

where the constant C_4 depends only on $\mathcal{L}$. From this we deduce (4.5) and $\|\bullet\| = \|\bullet\|_{L^2\to L^2}$. For instance,

$$\begin{aligned}\|[P,\chi_2]R(\lambda_0)\chi_1\|_{\mathrm{tr}} &\le C_4\|P[P,\chi_2]R(\lambda_0)\chi_1\| + C_4\|[P,\chi_2]R(\lambda_0)\chi_1\| \\ &= C_4\|[P,[P,\chi_2]R(\lambda_0)\chi_1\| + C_4(1+|\lambda_0|^2)\|[P,\chi_2]R(\lambda_0)\chi_1\| \\ &\le C_5.\end{aligned}$$

Here we used the facts that $\chi_2 \equiv 1$ on the support of χ_1, hence $[P,\chi_2]\chi_1 = 0$, and that $[P,[P,\chi_2]]$ $[P,\chi_2]$ are second and first order operators respectively and and that $R(\lambda_0)$ maps L^2 to $\mathcal{D}(P)$. The other terms in $K(\lambda,\lambda_0)$ are estimated similarly and that gives (4.5). (Finer estimates for large λ are possible – see [8, §4.3, §7.2] and [18]– but we concentrate here on uniformity near a given energy.)

Now, let $\Omega_3 = \{\lambda : |\lambda - \lambda_0| < R$ where R is large enough so that $\Omega_2 \subset \Omega_3$. It follows that for a constant C_3 depending only on Ω_3 and $\mathcal{L}$, (and hence only on Ω_2), we have

$$|\det(I + K(\lambda,\lambda_0)| \le e^{C_3}. \tag{4.7}$$

(For basic facts about determinants see for instance [8, §B.5].) Writing

$$(I + K(\lambda_0,\lambda_0))^{-1} = (I - (I + K(\lambda_0,\lambda_0))^{-1}K(\lambda_0,\lambda_0))$$

we obtain

$$\begin{aligned}|\det(I + K(\lambda_0,\lambda_0))|^{-1} &= |\det(I + K(\lambda_0,\lambda_0)^{-1}| \\ &\le \exp\left(\|(I + K(\lambda_0,\lambda_0))^{-1}\|\|K(\lambda_0,\lambda_0)\|_{\mathrm{tr}}\right) \le e^{C_4},\end{aligned}$$

that is

$$|\det(I + K(\lambda_0,\lambda_0))| \ge e^{-C_4}, \tag{4.8}$$

where C_4 depends only on λ_0 and $\mathcal{L}$. The Jensen formula (see for instance [19, §3.61]) then gives a bound on the number of zeros of $\det(I + K(\lambda,\lambda_0))$ in Ω_3. That proves the first bound in (4.1).

We can write

$$\det(I + K(\lambda_0,\lambda)) = e^{g(\lambda)} \prod_{\zeta\in\mathrm{Res}(P)\cap\Omega_3} (\lambda - \zeta),$$

where $g(\lambda)$ is holomorphic in Ω_3. From the upper bound (4.7) and the lower bound (4.8) we conclude that $|g(\lambda)| \le C_5$ in a smaller disc containing Ω_2, with C_5 depending only on the previous constants. (For instance we can use the Borel–Carathéodory inequality – see [19, §5.5].) Hence

$$|\det(I + K(\lambda_0,\lambda)| \ge e^{-C_6} \prod_{\zeta\in\mathrm{Res}(P)\cap\Omega_2} |\lambda - \zeta|, \quad \lambda \in \Omega_1,$$

To deduce the the second bound in (4.1) from this we use the inequality

$$\|(I + A)^{-1}\| \le \frac{\det(I + |A|)}{|\det(I + A)|}$$

which gives

$$\begin{aligned}\| \mathbb{1}_{r\leq R} R(\lambda) \mathbb{1}_{r\leq R} \| &= \| \mathbb{1}_{r\leq R} Q(\lambda,\lambda_0)\chi_3 (I+K(\lambda,\lambda_0))^{-1} \mathbb{1}_{r\leq R} \| \\ &\leq \| \mathbb{1}_{r\leq R} Q(\lambda,\lambda_0)\chi_3 \| |\det(I+|K(\lambda,\lambda_0))| |\det(I+K(\lambda,\lambda_0))|^{-1} \\ &\leq C_7 e^{\|K(\lambda,\lambda_0)\|_{\mathrm{tr}}} \prod_{\zeta\in \mathrm{Res}(P)\cap\Omega_2} |\lambda-\zeta|^{-1},\end{aligned}$$

for $\lambda\in\Omega_1$ and C_7 depending only on Ω_j's $\mathcal{L}$, and R. This completes the proof. □

Before proving Theorem 2 we will use the construction of the meromorphic continuation in the proof of Proposition 4.1 to give a general condition for smoothness of a family of resonances (see also [14]):

$$(P(t)-\lambda_0^2)^{-1}\in C^\infty((-t_0,t_0);\mathcal{L}(L^2,L^2)),\quad \operatorname{Im}\lambda_0>0. \tag{4.9}$$

That is the only property used in the proof of

Proposition 4.3. *Let $P(t)$ be the family of unbounded operators on L^2 (of a fixed metric graph) defined by (3.3). Let $R(\lambda,t)$ be the resolvent of $P(t)$ meromorphically continued to $\mathbb{C}$. Suppose that γ is a smooth Jordan curve such that $R(\lambda,t)$ has no poles on γ for $|t|<t_0$. Then for $\chi_j\in C^\infty_{\mathrm{c}}$, $j=1,2$,*

$$\int_\gamma \chi_1 R(\zeta,t)\chi_2 d\zeta \in C^\infty((-t_0,t_0);\mathcal{L}(L^2,L^2)). \tag{4.10}$$

In particular, if λ_0 is a simple pole of $R(\lambda,0)$ then there exist smooth families $t\mapsto\lambda(t)$ and $t\mapsto u(t)\in\mathcal{D}_{\mathrm{loc}}(P(t))$ such that $\lambda(0)=\lambda_0$, $\lambda(t)\in\mathrm{Res}(P(t))$ and $u(t)$ is a resonant state of $P(t)$ corresponding to $\lambda(t)$.

Proof. The proof of (4.10) under the condition (4.9) follows from (4.4) and the definitions of $Q(\lambda,\lambda_0)$ and $K(\lambda,\lambda_0)$. From that the conclusion about the deformation of a simple resonance is immediate – see [8, Theorems 4.7,4.9].

It remains to establish (4.9). Suppose $f\in L^2$ and define $u(t):=R(\lambda_0,t)f\in L^2$. Formally, $\dot u:=\partial_t u(t)$ satisfies (3.8) with $\dot z=0$ and $z=\lambda_0^2$. We can find a smooth family $g(t)\in L^2$ satisfying (3.9) with $u=u(t)$. We then have $\partial_t u(t)=g+R(\lambda_0,t)(-2\partial_t a(t)u(t)-G(t))$, where $G_m:=(-e^{-2a(t)}\partial_x^2-\lambda_0^2)g_m(t)$. By considering difference quotients a similar argument shows that $u(t)\in L^2$ is differentiable. The argument can be iterated showing that $u(t)\in C^\infty((-t_0,t_0),L^2)$ and that proves (4.9). □

We now give

Proof of Theorem 2. We proceed by contradiction by assuming that, for $0<\delta\ll\rho\ll 1>0$ to be chosen,

$$\mathrm{Res}(P)\cap(\Omega(\rho,\delta)+D(0,\delta))=\emptyset,\quad \Omega(\rho,\delta):=[\lambda_0-\rho,\lambda_0+\rho]-i[0,\delta]$$

does not contain any resonances. Choosing pre-compact open sets, independent of ε, ρ and δ, $\Omega(\rho,\delta) + D(0,\delta) \Subset \Omega_1 \Subset \Omega_2$ we apply Proposition 4.1 to see that for

$$\| \mathbb{1}_{r\le R}\, R(\lambda)\, \mathbb{1}_{r\le R} \| \le C_2 \delta^{-C_1}, \quad \lambda \in \Omega(\rho,\delta). \tag{4.11}$$

On the other hand, the resolvent estimate in the physical half-plane $\operatorname{Im}\lambda > 0$ and the fact that $\lambda_0 \in I \Subset (0,\infty)$, give

$$\| \mathbb{1}_{r\le R}\, R(\lambda)\, \mathbb{1}_{r\le R} \| \le C_3/\operatorname{Im}\lambda, \quad \operatorname{Im}\lambda > 0, \quad |\operatorname{Re}\lambda - \operatorname{Re}\lambda_0| < \rho. \tag{4.12}$$

To derive a contradiction we use the following simple lemma:

Lemma 4.4. *Suppose that $f(z)$ is holomorphic in a neighbourhood of $\Omega := [-\rho,\rho] + i[-\delta_-,\delta_+]$, $\delta_\pm > 0$. Suppose that, for $M > 1$, $M_\pm > 0$, and $0 < \delta_+ \le \delta_- < 1$,*

$$|f(z)| \le M_\pm, \quad \operatorname{Im} z = \pm\delta_+, \quad |\operatorname{Re} z| \le \rho, \qquad |f(z)| \le M, \quad z \in \Omega. \tag{4.13}$$

and that $\rho^2 > (1 + 2\log M)\delta_-^2$. Then

$$|f(0)| \le e M_+^{\theta} M_-^{1-\theta}, \quad \theta := \frac{\delta_-}{\delta_+ + \delta_-}. \tag{4.14}$$

Proof. We consider the following subharmonic function defined in a neighbourhood of Ω. To define it we put $m_\pm = \log M_\pm$, $m = \log M > 0$, $z = x + iy$, and

$$u(z) := \log|f(x+iy)| - \frac{\delta_- m_+ + \delta_+ m_- + y(m_+ - m_-)}{\delta_+ + \delta_-} - Kx^2 + Ky^2,$$

where $K := 2m/(\rho^2 - \delta_-^2)$. Then for $\operatorname{Im} z = \pm\delta_\pm$, $|\operatorname{Re} z| \le \rho$, $u(z) \le \delta_-^2 K \le 1$ since we assumed $\rho^2 > (1+2m)\delta_-^2$). When $|\operatorname{Re} z| = \rho$ then $u(z) \le 2m - K(\rho^2 - \delta_-^2) \le 0$. The maximum principle for subharmonic functions shows that $\log|f(0)| - \theta m_+ - (1-\theta)m_- \le 1$ and that concludes the proof. □

We apply this lemma to $f(z) := \langle \mathbb{1}_{r\le R}\, R(z+\lambda_0)\, \mathbb{1}_{r\le R}\, \varphi, \psi\rangle$, $\varphi, \psi \in L^2$, with $M_+ = C_3/\delta_+$, $M = M_- = C_2\delta^{-C_1}$. If we show that

$$|f(0)| \ll \frac{1}{\varepsilon}\|\varphi\|\|\psi\|, \tag{4.15}$$

we obtain a contradiction to (1.5) by putting $\psi = (P - \lambda_0^2)u$ and $\varphi = u$ and using the support property of u (the outgoing resolvent is the right inverse of $P - \lambda_0^2$ on compactly supported function):

$$1 = \langle R(\lambda_0)(P-\lambda_0^2)u, u\rangle = \langle \mathbb{1}_{r\le R}\, R(\lambda_0)\, \mathbb{1}_{r\le R}(P-\lambda_0)u, u\rangle \ll \frac{1}{\varepsilon}\varepsilon \ll 1.$$

For $\gamma < 1$ choose $\gamma < \gamma_1 < \gamma_2 < \gamma_3 < 1$ and put

$$\rho = \varepsilon^{\gamma_1}, \quad \delta_- = \varepsilon^{\gamma_2}, \quad \delta_+ = \varepsilon^{\gamma_3}.$$

Then (4.14) implies (4.15) and that completes the proof. □

For completeness we also include the following proposition which would be a converse to Theorem 2 for $\gamma = 1$. The more subtle higher dimensional case in the semiclassical setting was given by Stefanov [16].

Proposition 4.5. *Suppose that P satisfies the assumptions of Theorem 2 and let $R > 0, \delta > 0$. There exists a constant C_0 depending only of R, δ and $\mathcal{L}$ such that for any $0 < \varepsilon < \delta/2$,*

$$\begin{aligned} &D(\lambda_0, \varepsilon) \cap \operatorname{Res}(P) \neq \emptyset,\ \lambda_0 > \delta \implies \\ &\quad \exists\, u \in \mathcal{H}_R \cap \mathcal{D}_P,\ \|u\| = 1, \quad \|(P - \lambda_0^2)u\| \leq C_0 \varepsilon(\lambda_0 + \varepsilon). \end{aligned} \tag{4.16}$$

Proof. Suppose that λ a resonance of P with $|\lambda - \lambda_0| < \varepsilon$ and let v be the corresponding resonant state. Then in each infinite lead, $v_m(x) = a_m e^{i\lambda x}$, $1 \leq m \leq K$. As in (3.5),

$$\begin{aligned} \operatorname{Im}(\lambda^2)\|v\|^2_{\mathcal{H}_0} &= -\operatorname{Im} \sum_{m=1}^K \partial_x v_m(0)\bar{v}_m(0) = -\operatorname{Im} \sum_{m=1}^K i\lambda |a_m|^2 \\ &= -\operatorname{Re}\lambda \sum_{m=1}^K |a_m|^2, \end{aligned}$$

and since $\operatorname{Re}\lambda > \delta/2 > 0$, $\sum_{m=1}^K |a_m|^2 = 2|\operatorname{Im}\lambda| \|v\|_{\mathcal{H}_0} \leq 2\varepsilon \|v\|_{\mathcal{H}_0}$.

Suppose $r < R/2$ and $\chi \in C_{\mathrm{c}}^\infty([0,2)$ is equal to 1 on $[0,1]$. We then define $\widetilde{u} \in \mathcal{H}_R \cap \mathcal{D}_P$ by

$$\widetilde{u}_m(x) := \begin{cases} \chi(x/r)v_m(x), & 1 \leq m \leq K \\ v_m(x) & K+1 \leq m \leq K+M. \end{cases}$$

Now,

$$\|\widetilde{u}\|^2 = \|v\|^2_{\mathcal{H}_0} + \sum_{m=1}^K |a_m|^2 \int_{\mathbb{R}} e^{2|\operatorname{Im}\lambda| x} \chi(x/r)^2 dx = \|v\|^2_{\mathcal{H}_0}(1 + \mathcal{O}(\varepsilon r e^{2\varepsilon r})),$$

and hence,

$$\begin{aligned} \|(P - \lambda_0^2)\widetilde{u}\|^2 &= |\lambda^2 - \lambda_0^2|^2 \|\widetilde{u}\|^2 + \|[P, \chi(\bullet/r)]\widetilde{u}\|^2 \\ &\leq (2\varepsilon(\lambda_0 + \varepsilon))^2 \|\widetilde{u}\|^2 + C \sum_{m=1}^K |a_m|^2 (r^{-2} + (\lambda_0 + \varepsilon)^2) e^{2\varepsilon r} \\ &\leq C_{r,\delta} \varepsilon^2 (\lambda_0 + \varepsilon)^2 \|v\|^2_{\mathcal{H}_0}. \end{aligned}$$

We conclude that we can take $u := \tilde{u}/\|\tilde{u}\|$ as the quasimode. □

References

[1] G. Berkolaiko and P. Kuchment, *Introduction to Quantum Graphs*, Mathematical Surveys and Monographs, **186** AMS, 2013.

[2] G. Berkolaiko, P. Kuchment, *Dependence of the spectrum of a quantum graph on vertex conditions and edge lengths*, in *"Spectral Geometry"*, edited by A.H. Barnett, C.S. Gordon, P.A. Perry and A. Uribe. Proceedings of Symposia in Pure Mathematics **84**(2012).
[3] T. Carleman, *Über die asymptotische Verteilung der Eigenwerte partieller Differentialgleichungen,* Ber. Verk. Sächs. Akad. Wiss. Leipzig. Math.-Phys. Kl. **88**(1936), 119–134.
[4] Y. Colin de Verdière, *Pseudo-laplaciens. II,* Ann. Inst. Fourier **33**(1983), 87–113.
[5] H. Cornean, A. Jensen and G. Nenciu, *Metastable states when the Fermi golden rule constant vanishes*. Comm. Math. Phys. **334**(2015), 1189–1218.
[6] E.B. Davies, A. Pushnitski, *Non-Weyl resonance asymptotics for quantum graphs,* Analysis & PDE **4**(2012), 729–756.
[7] E.B. Davies, P. Exner, J. Lipovský, *Non-Weyl asymptotics for quantum graphs with general coupling conditions,* J. Phys. A **43**(2010), 474013.
[8] S. Dyatlov and M. Zworski, *Mathematical theory of scattering resonances,* book in preparation; http://math.mit.edu/~dyatlov/res/res.pdf
[9] P. Exner and J. Lipovský, *Resonances from perturbations of quantum graphs with rationally related edges,* J. Phys. A, **43**(2010), 105305.
[10] S. Gnutzmann, H. Schanz and U. Smilansky, *Topological resonances in scattering on networks (graphs),* Phys. Rev. Lett. **110**(2013), 094101.
[11] Y. Petridis and M. Risager, *Dissolving of cusp forms: higher-order Fermi's golden rules*, Mathematika **59**(2013), 269–301.
[12] R.S. Phillips and P. Sarnak, *Automorphic spectrum and Fermi's golden rule,* J. Anal. Math. **59**(1992), Festschrift on the occasion of the 70th birthday of Shmuel Agmon, 179–187.
[13] J. Sjöstrand and M. Zworski, *Complex scaling and the distribution of scattering poles,* J. Amer. Math. Soc. **4**(1991), 729–769.
[14] P. Stefanov, *Stability of resonances under smooth perturbations of the boundary,* Asymptotic Anal., **9**(1994), 291–296.
[15] P. Stefanov, *Quasimodes and resonances: sharp lower bounds,* Duke Math. J. **99**(1999), 75–92.
[16] P. Stefanov, *Resonances near the real axis imply existence of quasimodes,* C. R. Acad. Sci. Paris Sér. I Math., **330**(2000), 105–108.
[17] P. Stefanov and G. Vodev, *Neumann resonances in linear elasticity for an arbitrary body,* Comm. Math. Phys. **176**(1996), 645–659.
[18] S.H. Tang and M. Zworski, *From quasimodes to resonances,* Math. Res. Lett. **5**(1998), 261–272.
[19] E.C. Titchmarsh, *The theory of functions,* 2nd Edition, Oxford University Press, 1985.
[20] M. Zworski, *Distribution of poles for scattering on the real line,* J. Funct. Anal. **73**(1987), 277-296.

E-mail address: lee.minjae@math.berkeley.edu

E-mail address: zworski@math.berkeley.edu

Department of Mathematics, University of California, Berkeley, CA 94720, USA